

From Hawks and Doves to Self-Consistent Games of Territorial Behavior

Hanna Kokko,^{1,*} Andrés López-Sepulcre,^{1,2,†} and Lesley J. Morrell^{3,‡}

1. Laboratory of Ecological and Evolutionary Dynamics, Department of Biological and Environmental Science, University of Helsinki, P.O. Box 65 (Viikinkaari 1), FIN-00014 Helsinki, Finland;

2. Evolutionary Ecology Unit, Department of Biological and Environmental Science, University of Jyväskylä, P.O. Box 35, FIN-40014 Jyväskylä, Finland;

3. Division of Environmental and Evolutionary Biology, Institute of Biomedical and Life Sciences, University of Glasgow, Glasgow G12 8QQ, United Kingdom; and Institute of Integrative and Comparative Biology, University of Leeds, Leeds LS2 9JT, United Kingdom

Submitted August 26, 2005; Accepted February 15, 2006;
Electronically published May 11, 2006

Online enhancements: appendixes.

ABSTRACT: Explaining the “prior-residence effect” (automatic owner status of individuals who arrived first in an area) was one of the very first applications of game theory in animal behavior. These models, however, predict paradoxical solutions where intruders always win, with no satisfactory explanation for the absence of such cases in nature. We propose a solution based on new developments in evolutionary game theory. A self-consistent model with feedbacks between individual behavior and population dynamics produces qualitatively different frequency-dependent selection on intruders (floaters) than on territory owners. Starting with an ancestral population with no respect for ownership, the most likely evolutionary end point is complete or partial respect. Conventional rules of conflict resolution thus can rely on “uncorrelated asymmetries” without differences in resource-holding power or territory value, although they will be strengthened by such differences. We also review the empirical literature on animal contests, testing whether asymmetries in resource-holding power are required to explain the observations. Despite much empirical effort, results remain inconclusive, because experiments are often unable to distinguish between the motivation

of individuals to fight and the behavioral outcome of a contest. To help arrive at conclusive answers, we suggest a standardized empirical approach to quantify prior-residence effects.

Keywords: animal contests, ownership respect, prior-residence effect, self-consistent game theory, territoriality.

The animal kingdom provides countless examples of the “prior-residence effect,” the fact that individuals who arrived somewhere first appear to have a “psychological” advantage when competing over resources in that area. First-arrived individuals are more likely to win fights (some recent studies are Kemp and Wiklund 2004; Lefevre and Muehter 2004; Okada and Miyatake 2004; Switzer 2004; see table A1 in the online edition of the *American Naturalist* for a complete list), or alternatively, their ownership of the resources may go completely uncontested (e.g., Davies 1978; Baugh and Forester 1994; Turner 1994). Taxa in which prior-residence effects have been found range from sea urchins to mammals (see table A1), and the effect offers a powerful explanation of territorial contest resolution. In a situation where other individuals respect ownership, the owner can concentrate on more useful activities than resource defense, such as breeding.

But why should respect for ownership evolve? This question was one of the first applications of game theory to investigate animal behavior. Early hawk-dove games predicted that aggression (playing “hawk”) can be selected against, given sufficient costs of fighting (Maynard Smith and Price 1973). Applying the hawk-dove idea to territorial behavior requires defining a role asymmetry: one of the individuals is an owner, the other an intruder. Consequently, the list of strategies (hawk, dove) gains a new addition, the “bourgeois” strategy, in which an individual behaves aggressively only when in the owning role (Maynard Smith and Parker 1976). Indeed, the bourgeois strategy can be stable, and the outcome is termed an “uncorrelated asymmetry” (Maynard Smith and Parker 1976; Maynard Smith 1982): the difference between the players (owner vs. intruder) is sufficient to resolve the conflict despite the lack of correlation with fighting ability, value

* Corresponding author; e-mail: hanna.kokko@helsinki.fi.

† E-mail: andres.lopezsepulcre@helsinki.fi.

‡ E-mail: l.j.morrell@leeds.ac.uk.

of the resource to the individual, or any other effect that could bias the game in favor of one of the players. The uncorrelated-asymmetry hypothesis hence proposes that fights are settled based on purely conventional roles: it is cheaper for both players to adopt the convention that owners are respected simply because they are owners than to challenge the convention and end up in numerous fights as a consequence.

But the analysis of the ownership game reveals a strange feature, once one notices that the roles are mere arbitrary labels that allow settling the contest. There is nothing about the labels “owner” and “intruder” that make the owner the one more likely to adopt the role of the more aggressive player: the contest could be equally well settled by any other easily observable asymmetry, for example, “the player who is facing the sun plays dove.” To make matters worse, there are reasons why intruders in particular could be expected to be aggressive: if they cannot reproduce at all without a territory, they have very little to lose (Grafen 1987). For these reasons, it is essential to include the anti-bourgeois strategy, where owners always retreat (play “dove”) and intruders take their place (through playing hawk), in the set of possible strategies. The outcome is startling: antibourgeois is an evolutionarily stable strategy (ESS) exactly when bourgeois is too (Maynard Smith 1982; see table A2 in the online edition of the *American Naturalist*). Because the behavior predicted by this alternative appears counterintuitive, this solution is called the “paradoxical ESS,” as opposed to the “commonsense” nature of the bourgeois strategy (Maynard Smith and Parker 1976; Maynard Smith 1982).

The current consensus is that the uncorrelated-asymmetry hypothesis does not perform well as an explanation of territorial behavior (e.g., Hardy 1998). This is not only because of the strange duality it predicts between paradoxical and commonsense strategies. When looking closely, one usually finds some feature of individuals that is consistently different between resource owners and intruders, be it resource-holding power (RHP; e.g., Lindström 1992; Lozano 1994; Kemp and Wiklund 2001; Pryke and Andersson 2003)—which sometimes is causally enhanced through ownership, for example, when basking butterflies warm up in the sun (Stutt and Willmer 1998; but see Kemp and Wiklund 2004)—or the value of a territory, which again can increase with time spent as a territory owner (as argued by, e.g., Krebs [1982]; Tobias [1997]; Stokkebo and Hardy [2000]). This means that one does not have to rely on uncorrelated asymmetries as a solution.

On the other hand, some theoretical work aims to show that uncorrelated asymmetries do not necessarily predict this strange duality, but certain mechanisms ensure that the commonsense strategy prevails, for example, repeated interactions (Mesterton-Gibbons 1992), errors in decision

making (Morrell and Kokko 2003), or physiological correlations between ways of responding to draws versus losses (Morrell and Kokko 2005). However, a closer look reveals that these cases always include some a priori assumption that makes owners more likely to win fights: Mesterton-Gibbons (1992) predefines a probability $>1/2$ that owners win, Morrell and Kokko (2003) assume that errors make fight outcomes correlate with true fighting ability, and the physiological correlations assumed by Morrell and Kokko (2005) have a similar effect. Thus, one could conclude—as, for example, Mesterton-Gibbons (1992) has done—that ownership as a pure convention does not work, in the sense of being able to predict commonsense solutions of ownership respect; some correlated asymmetries are always needed, perhaps together with cognitive mechanisms that allow assessment of the opponent’s strength (Parker and Rubenstein 1981).

Here our aim is to challenge this consensus, based on theoretical considerations as well as an investigation of the way empirical tests are conducted. Regarding theoretical developments, we will show that game theory can, after all, produce models based on uncorrelated asymmetries that produce respect for ownership. The key is to make models self-consistent (Houston et al. 2005; Houston and McNamara 2006); self-consistent models are also called “ecogenetic” models (Eshel and Sansone 1995) or models with eco-evolutionary feedback (Le Galliard et al. 2005). The rationale is as follows. It is not logically consistent to analyze a game where the value of a territory and costs of fighting are constants V and C , respectively. For example, if fights are frequent, the value of a territory must necessarily decline, because challenges occur more often (Eshel and Sansone 1995; Houston and McNamara 2006). Likewise, the same costs of fighting must lead to a shorter life span if fights are very frequent; and if mortality is high in the population as a whole, a nonaggressive strategy that simply waits for a vacancy must perform relatively better, since territorial turnover is high.

By explicitly investigating all the options awaiting individuals throughout their lifetime, our models are made self-consistent, but they also incorporate Maynard Smith’s (1982, p. 96) idea of “infinite regress.” This argument has been used to explain away paradoxical strategies: if such a strategy really did prevail, there would be no time left for breeding because population members keep moving around, finding new places to settle only to be soon replaced by another individual. However, the exact consequences of an infinite regress have never been formally quantified; thus, its ability to remove paradoxical solutions has not been put under test. Here we provide a model that allows an infinite regress to take place (within the limits of the lifetime of individuals and their mobility) and calculates the consequent reproductive rates, using the as-

sumption that individuals can reproduce until they are displaced by another. After developing self-consistent models of territory ownership with and without resource-holding power differences, we comment on the empirical literature on correlated and uncorrelated asymmetries and give recommendations for future research.

Modeling Territorial Behavior with Population-Level Feedbacks

Our aim is to provide a self-consistent account of the aggressiveness of owners and intruders and to see if we can predict respect for ownership such that intruders are less aggressive than owners upon encounter. Grafen (1987) pointed out that when predicting the behavior of an intruder, one should consider how “desperate” it might be: what are the options available if it does not fight for the contested resource? However, Grafen’s argument does not immediately predict a difference in the aggression of owners and intruders. Being desperate merely increases the payoff difference between having or not having the resource, but this applies to owners who might lose the resource as well as to intruders who might gain it (table A2).

However, taking into account population dynamics could change the situation. A nonbreeding intruder (often called a “floater”; Zack and Stutchbury 1992) will find itself in a more relaxed situation if territory owners often succumb to death: simply waiting for a vacancy to arise might then be enough. This must be evaluated against the floater’s own survival prospects during the appropriate time. More important still, variables such as the rates of finding vacancies will not be fixed for each species but will themselves evolve; for example, if many floaters kill themselves by fighting aggressively, it might become relatively better to wait, because the numbers of competitors will be lower.

Here we examine the dynamics of floating and breeding with population-level feedback. After solving the model with no RHP differences, we introduce variation in RHP, to see how differences in fighting ability influence the distribution of commonsense versus paradoxical outcomes. This allows us to examine theoretically the hypothesis that ownership is respected because strong individuals tend to accumulate as owners (Parker 1974; Leimar and Enquist 1984; Alcock and Bailey 1997; Kemp and Wiklund 2004).

Evolutionarily Stable Behavior with No Differences in RHP

We begin the description of the model by defining the possible strategies. We consider a population consisting of breeders (territorial owners) and nonterritorial floaters. If these meet (according to rules defined below), the owner and floater can each play “daring” or “careful.” These

correspond to hawks and doves in older models. The strategy is described by the probabilities that an individual plays daring if it is a territorial owner or a floater. For brevity, we call the probability of playing daring the “aggressiveness” of an individual, denoted x for owners and y for floaters. A daring floater is an intruder willing to fight, while a daring owner defends his territory aggressively. If only one individual of the two plays daring, it takes over the territory. If neither is daring, either player obtains the territory with probability 1/2, and the other one leaves. If both are daring, a fight ensues. In this section, we assume no RHP asymmetry, so that one of the players (the winner) obtains the territory and the other one (the loser) leaves, again with probabilities 1/2.

We now define the parameters that are fixed for a population (Greek letters in table A3 in the online edition of the *American Naturalist*). We assume that breeding occurs continuously. Time units are scaled such that territory owners produce one offspring per time unit. Territory owners have a background mortality rate of μ_T ; this rate excludes deaths associated with fights. Because owners produce offspring at a rate 1 and floaters do not produce offspring, $\mu_T < 1$ will be necessary to avoid population extinction. Newborn individuals become nonbreeding floaters, who can then either fill territories vacated by the death of the territory owner or obtain territories by usurping the current owner. The background mortality of floaters (excluding deaths associated with fights) equals μ_F .

If an owner dies, the nearest floater takes over the vacancy. We assume that in addition to encountering possible vacant territories, floaters sample θ occupied territories per time unit and that they meet the territory owner every time they sample such a territory. If both individuals then play daring, a fight ensues.

Fights can be lethal for the loser, who will die as a result of the fight with probability δ . The background mortality rates μ_T and μ_F , the lethal injury probability δ , and the rate of territory inspection θ are the only parameters that take fixed values for each population. There are additional population parameters whose values depend on the behavior of population members; we call these “feedback parameters.” Such parameters are the number of floaters per territory owner (denoted n) and the total mortality rates of territory owners (m_T) and floaters (m_F). When necessary, we will use notations such as $n(x, y)$ to indicate dependence on population strategies. For brevity, however, we usually use the shorter notation n . To make it clear which parameters are fixed in a population and which ones evolve with the strategy played, we have used Greek letters to denote fixed values and Latin characters for parameters and variables that respond to the strategy in use in the population (see table A3 for a list of notation).

The total mortality rate of owners increases with the

relative number of potential intruders n and their rate of territory inspection θ , because intruders cause fights that can be lethal (with probability δ). Because an inspection causes a fight with probability xy and the probability of losing is $1/2$, the total mortality of owners equals

$$m_T = \mu_T + \frac{xy n \theta \delta}{2}. \quad (1)$$

The total mortality rate of floaters, m_F , is derived similarly, but the rate of territory inspection is θ per floater; thus,

$$m_F = \mu_F + \frac{xy \theta \delta}{2}. \quad (2)$$

The number of floaters per territory, n , increases at a rate of 1 per time unit through births and decreases at a rate $m_F n$ through deaths of floaters and at a rate m_T because of floaters either finding vacancies created by the death of the territorial or taking over territories of owners who died in the contest. Note that floaters can become territorials through usurpation of an owner, but in cases of no mortality, this does not change the net number of floaters, and the associated deaths are taken into account in the total mortality rates. At equilibrium, the number of floaters, n , does not change over time:

$$\frac{dn}{dt} = 1 - m_F n - m_T = 0. \quad (3)$$

This gives the equilibrium number of floaters per territory,

$$n = \frac{1 - m_T}{m_F} = \frac{1 - \mu_T}{\mu_F + \theta xy \delta}. \quad (4)$$

The rate at which an individual floater finds and occupies vacant territories is again a feedback parameter,

$$v = \frac{\mu_T}{n}. \quad (5)$$

Evolutionary Equilibria and Their Stability Conditions. We can derive the following results (the derivation and the exact values for all criteria are detailed in app. B in the online edition of the *American Naturalist*):

1. No ESS exists where owners play a mixed strategy ($0 < x^* < 1$).

2. The only possible ESS of the form $\{0, y^*\}$ is the pure paradoxical strategy $\{0, 1\}$.

From results 1 and 2, it follows that the only possible equilibria are the paradoxical equilibrium $\{0, 1\}$, a “no respect” equilibrium (i.e., all individuals always fight) $\{1,$

$1\}$, the commonsense “complete-respect” equilibrium $\{1, 0\}$, and a semimixed “partial-respect” strategy where owners always defend, and floaters sometimes challenge them: $\{1, y^*\}$ with $0 < y^* < 1$. While the partial-respect strategy does not show full respect for ownership, it still has commonsense features: owners always defend their property, and intruders at least sometimes retreat without challenging them. It is noteworthy that the “hippie world” solution $\{0, 0\}$, where resources are shared without aggression, is never stable.

3. The commonsense complete-respect solution, where owners defend their territories and intruders respect ownership, is an ESS when the background mortality of territory owners is large and fighting is risky.

4. The paradoxical solution, where owners retreat and intruders replace them without fights, is an ESS when the background mortality of breeders is high, that of floaters is low, the number of territories they can inspect per unit time θ is high, and fight costs are high. In this case, reproductive values of owners and floaters do not differ from each other hugely: an owner does not lose as much from stepping down as it would under different circumstances.

5. As in solution 3, the no-respect solution, where individuals always fight when meeting each other ($x = y = 1$), is an ESS when the background mortality of floaters is high, breeder mortality is low, and floaters do not encounter new territories quickly. These conditions imply that floaters are “desperadoes”: breeding possibilities are rarely encountered, and floaters are short-lived and thus in a hurry. This equilibrium also becomes more likely if fight costs δ are low.

6. The semimixed partial-respect ESS with $x^* = 1$, $0 < y^* < 1$ can be stable, and the value of y^* is given in appendix B. The degree of respect increases (i.e., floater aggression decreases) with the costs of fighting, the rate of territory encounter, and the background mortality of breeders. When the mixed equilibrium exists, it always coexists with the paradoxical equilibrium.

We also show (app. B) that the conditions that stabilize the commonsense equilibrium are also sufficient to make the paradoxical ESS exist. The converse is not true.

Evolutionary Trajectories with or without Variation in RHP. The previous section showed that full or partial respect can be evolutionarily stable, but such solutions always coexist with the paradoxical equilibrium: introducing population feedback does not destroy the paradoxical strategy. The paradoxical strategy, on the other hand, can exist without coexisting with a “respectful” strategy. This seems to suggest that population feedback is of little use when explaining the prevalence of respect for ownership in nature. However, it is clearly of interest to ask which ESS is reached from various starting conditions: the basins

of attraction—that is, the regions of initial values (x, y) from which evolution proceeds to each equilibrium—are not necessarily equally large for the various ESSs. One must additionally evaluate how biologically likely each starting point $\{x, y\}$ is for the initial evolution of territoriality. It makes little sense to consider starting values that can never be stable and are thus unlikely to be reached at any point during evolution (such as hippie world, $\{0, 0\}$).

Our numerical procedure for calculating evolutionary trajectories is outlined in appendix B. In this section, we also provide an extension of the ESS analysis by relaxing the assumption that there are no differences in individuals' fighting abilities. The possibility that high-RHP individuals tend to accumulate as owners was already proposed by Parker (1974) to explain ownership respect, but to our knowledge it has not been properly developed theoretically. In that case, residency itself will convey information about relative RHP. We include this in our model by assuming that individuals can fall into two categories: they are either strong or weak fighters. This trait is randomly determined at birth (we assume no heritability of the character), with a proportion π of individuals born strong.

Because we are interested in whether differences in fighting ability can lead to general respect for ownership such that ownership functions as the sole cue that settles conflicts, we assume that individuals know only their status as floater or owner. The status can, of course, correlate with fighting ability, and the evolutionary success of strategies is calculated accordingly. In contests between two individuals of the same fighting ability, the fight leads to victory or loss and possible death as in the previous section. If a strong and a weak individual fight, the strong one wins with probability $(1 + \alpha)/2$, where α describes fight predictability, that is, the degree to which strength gives an advantage in fights. If $\alpha = 0$, outcomes are randomly determined, and the model then becomes identical to the “no-RHP-difference” model developed in the previous section, whereas with $\alpha = 1$, the stronger contestant invariably wins.

When $\alpha = 0$ (i.e., no RHP differences), the numerical procedure shows that the various ESSs can be reached by evolution (fig. 1). If one considers that the most relevant starting point for an ancestral population is one in which respect for ownership has not yet evolved and resources are always contested (no respect; $x = y = 1$ in the upper right corner of both panels in fig. 1), evolution either stays at no respect (fig. 1A) or proceeds to the equilibrium with partial respect of ownership rather than the paradoxical equilibrium.

Figure 1 shows only the solutions for one particular choice of values of population parameters (mortalities and efficiency of territorial sampling). However, its results are not an isolated example. To obtain an overview of how commonly evolution can lead to paradoxical versus com-

monsense strategies when starting from no respect, we checked 1,000 random choices for parameter values: parameter values were randomly picked from an even distribution that ranged between 0 and 1 for the parameters μ_T , μ_F , and δ and between 0 and 10 for θ . Then we calculated the evolutionary trajectory, starting from $x = 1$, $y = 1$. In 398 cases, the population stayed at the no-respect equilibrium. In 592 cases, the evolutionary end point was commonsense or partially commonsense, and in a mere 10 cases, the paradoxical solution evolved. In these 10 paradoxical cases, mean floater mortality was only 0.0153, and the baseline mortality of owners was, on average, 44 times that of floaters (range 1.43–114.6). We may conclude that evolution can proceed from no respect toward the paradoxical ESS, but this appears to require an extremely unlikely mortality pattern where territory owners suffer vastly higher mortality than floaters.

Figure 2A depicts a typical result in the absence of RHP differences, while figure 2B (with $\alpha = 1$) adds these differences. It is clear that differences in fighting ability make intruders more careful: because strong owners have accumulated in the territorial population and weak fighters have often lost fights, floaters are more likely to be weak than owners. At the equilibrium $x^* = 1$ and $y^* = 0.23$, 94% of owners are strong, compared with 48% of floaters (calculations follow app. B). Floaters therefore enjoy smaller expected payoffs from engaging in a fight than they do in figure 2A, which explains their greater reluctance. However, the difference between figure 2A and figure 2B is quantitative rather than qualitative: in both cases, evolution proceeds toward a partial-respect equilibrium, and the only difference is in the value of y^* .

How powerful are RHP differences in producing qualitatively different outcomes? For example, can they switch the territorial system from the no-respect state to one with respect for ownership, either partial or complete? Figure 3 shows the effect of fight predictability α on the evolutionary end point when evolution begins in the no-respect state. In all cases of figure 3, owners always defend their property, and increasing α makes intruders less aggressive (i.e., decreases y^*). This includes qualitative changes; for example, when $\delta = 0.35$, no respect changes to partial respect, with $y^* = 0.78$, while α increases from 0 to 1 (fig. 3).

Nevertheless, partial respect in the above example is not very strong—78% of intruders still challenged the owner. Overall, figure 3 shows that the influence of RHP differences is fairly slight. Solutions are never paradoxical in figure 3, even if we assume no RHP differences (at $\alpha = 0$), and the lines delimiting different solutions are vertical or close to vertical in most parts of figure 3, which indicates that predictability of fight outcomes, α , merely slightly shifts regions of different solutions toward less frequent challenges by intruders; in other words, RHP differences




Figure 1: Evolutionary trajectories of the feedback model without RHP differences. Dots mark stable equilibria. Dotted lines describe evolution toward the paradoxical strategy, solid lines evolution toward any other kind. *A*, Fighting carries fairly low mortality risk ($\delta = 0.1$), leading to two evolutionary end points: the paradoxical strategy or the no-respect equilibrium. Other parameters: $\theta = 10$, $\mu_T = \mu_F = 0.5$. *B*, Fighting carries high mortality risk ($\delta = 0.5$), which makes no respect unstable and partial respect stable. Other parameters are as in *A*.




Figure 2: Comparison of model with and without RHP differences. *A*, No RHP differences and parameters $\theta = 10$, $\mu_T = 1.0$, and $\mu_F = 1.2$ lead to partial respect being stable in a similar vein as in figure 1. At this equilibrium, intruders challenge owners in approximately 40% of encounters. *B*, Assuming RHP differences with $\alpha = 1$, $\pi = 0.5$ leads to similar evolutionary trajectories, but there is now more respect: intruders challenge owners less often, in approximately 23% of encounters, than in *A*. Other parameters are as in *A*.




Figure 3: Evolutionary end points when evolution starts in the no-respect state ($x = y = 1$). Types of solutions are derived with varying values of fight predictability α , as indicated on the y -axis, and varying values of (A) fighting cost δ , (B) territory encounter rate θ , (C) breeder mortality μ_T , and (D) floater mortality μ_F , as indicated on each x -axis. Parameters, where not varied, are $\delta = 0.4$, $\theta = 5$, $\mu_F = \mu_T = 0.1$, and $\pi = 0.5$. The graph divides the parameter space into regions: “NR” indicates that the evolutionary end point is no respect, $x^* = y^* = 1$, and “CR” that it is complete respect, $x^* = 1, y^* = 0$. In any other region, the solution features partial respect, with the numbers indicating contour lines for the evolutionarily stable aggressiveness of intruders, y^* .

introduce a slightly stronger tendency to respect ownership. The tendency per se was already introduced by the feedback model with no RHP differences.

Discussion

The prior-residence effect was first reported by Davies (1978), who observed territorial contests in speckled wood butterflies *Pararge aegeria*. Since then, it has been found in a multitude of species (see table A1 for examples), and we are aware of only two species in which the paradoxical solution (i.e., intruders attack and residents retreat systematically) has been observed: a social

spider *Oecobius civitas* (Burgess 1976; but see Mesterton-Gibbons and Adams 1998) and the captive goldeye fish *Hiodon alosoides* (Fernet and Smith 1976). Clearly, this means that game theoretical models of animal contests cannot be taken seriously unless they are able to predict the extreme rarity of paradoxical solutions. Previous solutions have relied on assumptions such as cognitive capabilities allowing RHP assessment (Parker and Rubenstein 1981). Our reanalysis of old hawk-dove games, using self-consistent modeling of population processes, explains why paradoxical solutions are unlikely to develop in nature, regardless of how developed the cognitive machinery is. Frequency dependence operates differently

for intruders and for owners, and this has both theoretical and empirical consequences.

Regarding theoretical advances of applications of game theory in animal behavior, it is increasingly being realized that payoffs cannot be considered fixed parameters: when behavior evolves, not only does the expected behavior of a conspecific individual change, but the numbers of conspecific competitors, mates, etc., typically change, too (Le Galliard et al. 2005; López-Sepulcre and Kokko 2005; Houston and McNamara 2006). In other words, the ecological setting in which a population finds itself depends on the strategies used in the population. Here we have applied self-consistency to the very first problem—aggression and resource defense—where game-theoretic logic was introduced to the study of animal behavior.

Eshel and Sansone (1995) argue that fights lead to owners and intruders swapping roles in a way that maintains the arbitrary nature of these roles and that paradoxical solutions can thus prevail. This echoes the argument of our table A2, where both intruders and owners compare the same values of winning, V , and getting injured in a fight, C ; hence, they experience the same payoffs. This logic does not hold, however, when population feedback is taken into account: it changes the payoffs of the game as aggressiveness levels change, mediated via population density and the number of vacancies present.

The consequences differ for floaters and owners. The best option for a floater can change from “challenge” to “do not challenge” when floaters become more aggressive: more fights in the population mean that fellow floaters die more often, thus shortening the wait to find an available territorial vacancy. The less risky option of non-aggressive waiting therefore becomes better, creating negative frequency dependence on the floater strategy and stabilizing a mixed strategy for floaters. (The argument would become stronger still had we allowed owners to die after the fight even if they won it; increased aggressiveness would then imply more vacancies created this way.) Owners, on the other hand, do not experience negative frequency dependence: they should never switch from “defend” to “do not defend” when daring owners become more prevalent in the population. More fighting means that the number of floaters decreases, which makes it more, rather than less, beneficial for owners to be aggressive: the value of a territory that is rarely challenged is higher and should be defended more.

Population feedback can therefore generate owner-intruder asymmetries in behavior without a need to assume that owners are a priori more likely to win. This improves the realism of game theory models, as does the fact that we often predict partial respect, where breeders usually enjoy ownership rights but these are sometimes contested and takeovers may take place. Complete respect

begs the question of why intruders should intrude at all if they always retreat (Grafen 1987), a problem not present in solutions featuring partial respect.

To empiricists, our most important message is that evidence for asymmetries between owners and intruders does not rule out that an underlying uncorrelated asymmetry has a strong influence on ownership evolution. The prevailing interpretation is that the prior-residence effect is not a sign of a pure convention, as proposed by Maynard Smith and Parker (1976). Rather, it is viewed a consequence of residence being correlated to asymmetries in competitive ability or value of the resource (e.g., Stokkebo and Hardy 2000). It has been suggested (Kemp and Wiklund 2001) that the role of conventions must be minor, because careful examination almost always reveals some asymmetry between owners and intruders in either RHP or the value of the territory. Together with the possibility of RHP assessment (Parker and Rubenstein 1981), asymmetries then appear sufficient to explain the owner-intruder asymmetry in strategies. For example, Kemp and Wiklund (2004) reexamined prior-residence effects in the speckled wood butterfly (the same species used by Davies [1978]) in a carefully controlled experimental setting. Through a clever series of experimentation, they found evidence supporting the idea that more intrinsically aggressive males accumulate as residents, which could then explain the prior-residence effect, too.

Our figure 3 suggests that RHP differences can play some role: solutions can incorporate differences in fighting ability and the accumulation of strong owners, and this strengthens ownership respect to some extent. Yet this does not mean that the signal of the underlying convention is absent, since predictions regarding ownership respect remain qualitatively similar if RHP differences diminish and vanish (fig. 3). Certainly, the results of controlled experiments such as those by Kemp and Wiklund (2004) are not compatible with partial- or complete-respect solutions in our model without RHP differences. However, we also predict that no respect can prevail as a stable solution, and, perhaps surprisingly, the results of Kemp and Wiklund can be completely reconciled with our approach if they are described as a no-respect solution with RHP differences. This solution is found in figure 3 under a wide set of parameter values. Under such a scenario, we would predict no difference in the willingness to initiate fights (which can be termed the “motivation” to fight, *sensu* Härdling et al. 2004), while the observed fight outcomes can be biased toward owners winning if strong individuals have accumulated in the owner population. Kemp and Wiklund (2004) reported that contests always ensued immediately; thus, their results regarding motivation are consistent with this interpretation.

This difference between fight initiation and fight out-

come highlights a large gap between empirical and theoretical approaches. Theoretical studies typically model the evolution of fighting strategies (motivational aspects, i.e., whether to attack or retreat), while most empirical studies record the outcome of those fights (i.e., who is observed to win). Although the outcome of a fight will be determined partly by the strategy (motivation) of both opponents (and some studies show a correlation between them; Grossman 1980; Mayr and Berger 1992; Chellappa et al. 1999; Maan et al. 2001), observing only fight outcomes is not sufficient to distinguish between strategies. For example, intruders are sometimes predicted to fight even when their chances to win are low (Grafen 1987). If one were to measure contest outcome in such a system, residents would win most of the time, suggesting a prior-residence effect, even if the intruder's strategy showed no respect for ownership at all.

While it is simple to define strategies of "intention" or motivation in theoretical models, it is far from easy to detect them in nature. If conflicts are solved by the convention "resident wins," most of those interactions are likely to be missed, since floaters end up not challenging resource owners at all (Grafen 1987). In fact, once a fight is observed, one can interpret that it occurred because both resident and floater have played daringly, and the task of the empiricist is then to try to assess the fraction of encounters that led to such a challenge. Most empirical studies, however, focus only on realized fights. For example, in a study investigating fighting tactics in mouthbrooding cichlid fish *Oreochromis mossambicus*, Turner (1994) excluded from the analyses all cases where fights did not occur because of intruders retreating—which added up to 95% of the total cases. Clearly, the respect of ownership becomes much more impressive if such cases are included. Other studies have applied similar selection criteria (e.g., Harvey and Corbet 1986; Bridge et al. 2000). A focus only on realized fights leads to a dilution of the importance of respect for prior ownership, which gives another reason why the role of conventions may remain underappreciated. Naturally, empiricists need to be aware of possible misassessments of the opposite kind, too: not all intrusions that end up in the intruder fleeing were meant to be territorial challenges in the first place (Grafen 1987).

We appreciate, of course, the complications that arise when trying to detect the historical evolutionary pathway. RHP differences or value asymmetries exist in many systems, often together with an appropriate assessment mechanism (Parker and Rubenstein 1981), and detecting "which came first" may require phylogenetically controlled comparative approaches. Most studies where residency is tested against RHP conclude that competitive ability (RHP) is a stronger determinant of fight outcome than residency. Evidence supporting strong prior-residence effects is most

often found in three scenarios (see table A1 for a more detailed list of studies): first, when individuals are more or less matched in size (e.g., Hammerstein and Riechert 1988; Rosenberg and Enquist 1991; Stuart-Smith and Boutin 1994; Beaugrand et al. 1996; Petersen and Hardy 1996; López and Martín 2001); second, when individuals with a higher RHP tend to accumulate as residents (e.g., Lindström 1992; Alcock and Bailey 1997; Pratt et al. 2003; Pryke and Andersson 2003); and third, when residence confers a higher RHP—as is the case with resident butterflies being warmer (Stutt and Willmer 1998) or in organisms that defend burrows being in a better position to fight (e.g., Magnhagen and Kvarnemo 1989; Ranta and Lindström 1993; Jennions and Backwell 1996). This is in keeping with our model, where RHP differences simply strengthen a convention that already exists. In appendix B, we outline an empirical method to quantify the effects of asymmetries in the presence of an underlying convention.

It also remains to be tested whether conventions really do arise through frequency dependence in the payoff structure, as assumed by our model. This poses obvious challenges: to study population-level feedback, one requires, ideally, much more extensive information about the nature of density-dependent population regulation than is usually available (e.g., Kokko and Rankin 2006). However, differences in payoffs have been shown to correspond to changes in individual behavior (Hammerstein and Riechert 1988), which suggests that such an approach could prove extremely fruitful.

Further biological complexities certainly have the potential to influence the fitness prospects of a floater: learning can play a role in the development of ownership (Stamps and Krishnan 1999, 2001; Morrell and Kokko 2003) as well as the possibility that intruders negotiate some space for themselves by squeezing in between existing territories (Stamps and Krishnan 2001; Pereira et al. 2003; López-Sepulcre and Kokko 2005). Regarding such additions, our modeling simply shows that a potentially fundamental biological process strongly predisposes species to exhibit a prior-residence effect, even if no additional respect-strengthening mechanisms (such as RHP assessment; Parker and Rubenstein 1981) exist in a particular case. Together with the overwhelming ubiquity of the prior-residence effect regardless of the nature of the resource, type of weaponry, and other biological details of the diverse taxa in which such effects are reported (see table A1), this suggests to us that the prior-residence effect reflects a more fundamental biological process than a mere assertion that some asymmetry or other is usually present in animal populations.

Acknowledgments

We thank I. Hardy, D. Kemp, and two anonymous reviewers for useful comments on the manuscript; L. Frank Baum for inspiration; and P. Hammerstein and J. McNamara for discussions. Funding was provided by the Academy of Finland and by the Natural Environment Research Council.

Literature Cited

- Alcock, J., and W. J. Bailey. 1997. Success in territorial defence by male tarantula hawk wasps *Hemipepsis ustulata*: the role of residency. *Ecological Entomology* 22:377–383.
- Baugh, J. R., and D. C. Forester. 1994. Prior residence effect in the dart-poison frog, *Dendrobates pumilio*. *Behaviour* 131:207–224.
- Beaugrand, J. P., D. Payette, and C. Goulet. 1996. Conflict outcome in male green swordtail fish dyads (*Xiphophorus helleri*): interaction of body size, prior dominance/subordination experience and prior residency. *Behaviour* 133:303–319.
- Bridge, A. P., R. W. Elwood, and J. T. A. Dick. 2000. Imperfect assessment and limited information preclude optimal strategies in male-male fights in the orb-weaving spider *Metellina megei*. *Proceedings of the Royal Society of London B* 267:273–279.
- Burgess, J. W. 1976. Social spiders. *Scientific American* 234:100–106.
- Chellappa, S., M. E. Yamamoto, M. Cacho, and F. A. Huntingford. 1999. Prior residence, body size and the dynamics of territorial disputes between male freshwater angelfish. *Journal of Fish Biology* 55:1163–1170.
- Davies, N. B. 1978. Territorial defence in the speckled wood butterfly (*Pararge aegeria*): the resident always wins. *Animal Behaviour* 26:138–147.
- Eshel, I., and E. Sansone. 1995. Owner-intruder conflict, Grafen effect and self-assessment: the bourgeois principle re-examined. *Journal of Theoretical Biology* 177:341–356.
- Fernet, D. A., and R. J. F. Smith. 1976. Agonistic behavior of captive goldfeye (*Hiodon alosoides*). *Journal of the Fisheries Research Board of Canada* 33:122–127.
- Grafen, A. 1987. The logic of divisively asymmetric contests: respect for ownership and the desperado effect. *Animal Behaviour* 35:462–467.
- Grossman, G. D. 1980. Food, fights, and burrows: the adaptive significance of intraspecific aggression in the bay goby (Pisces: Gobiidae). *Oecologia (Berlin)* 45:261–266.
- Hammerstein, P., and S. E. Riechert. 1988. Payoffs and strategies in territorial contests: ESS analyses of two ecotypes of the spider *Agelenopsis aperta*. *Evolutionary Ecology* 2:115–138.
- Hårdling, R., H. Kokko, and R. W. Elwood. 2004. Priority versus brute force: when should males begin guarding resources? *American Naturalist* 163:240–252.
- Hardy, I. C. W. 1998. Butterfly battles: on conventional contests and hot property. *Trends in Ecology & Evolution* 13:385–386.
- Harvey, I. F., and P. S. Corbet. 1986. Territorial interactions between larvae of the dragonfly *Pyrrosoma nymphula*: outcome of encounters. *Animal Behaviour* 34:1550–1561.
- Houston, A. I., and J. M. McNamara. 2006. John Maynard Smith and the importance of consistency in evolutionary game theory. *Biology and Philosophy (forthcoming)*, doi: 10.1007/s10539-005-9016-4.
- Houston, A. I., T. Szekely, and J. M. McNamara. 2005. Conflict between parents over care. *Trends in Ecology & Evolution* 20:33–38.
- Jennions, M. D., and P. R. Y. Backwell. 1996. Residency and size affect fight duration and outcome in the fiddler crab *Uca annulipes*. *Biological Journal of the Linnean Society* 57:293–306.
- Kemp, D. J., and C. Wiklund. 2001. Fighting without weaponry: a review of male-male contest competition in butterflies. *Behavioral Ecology and Sociobiology* 49:429–442.
- . 2004. Residency effects in animal contests. *Proceedings of the Royal Society of London B* 271:1707–1711.
- Kokko, H., and D. J. Rankin. 2006. Lonely hearts or sex in the city? density-dependent effects in mating systems. *Philosophical Transactions of the Royal Society of London B* 361:319–334.
- Krebs, J. R. 1982. Territorial defence in the great tit (*Parus major*): do residents always win? *Behavioral Ecology and Sociobiology* 11:185–194.
- Lefevre, K. L., and V. R. Muehler. 2004. Competition for mating resources in a territorial damselfly (Odonata: Calopterygidae). *Studies on Neotropical Fauna and Environment* 39:159–165.
- Le Galliard, J. F., R. Ferrière, and U. Dieckmann. 2005. Adaptive evolution of social traits: origin, trajectories, and correlations of altruism and mobility. *American Naturalist* 165:206–224.
- Leimar, O., and M. Enquist. 1984. Effects of asymmetries in owner intruder conflicts. *Journal of Theoretical Biology* 111:475–491.
- Lindström, K. 1992. The effect of resource holding potential, nest size and information about resource quality on the outcome of intruder-owner conflicts in the sand goby. *Behavioral Ecology and Sociobiology* 30:53–58.
- López, P., and J. Martín. 2001. Fighting rules and rival recognition reduce costs of aggression in male lizards *Podarcis hispanica*. *Behavioral Ecology and Sociobiology* 49:111–116.
- López-Sepulcre, A., and H. Kokko. 2005. Territorial defense, territory size, and population regulation. *American Naturalist* 166:317–329.
- Lozano, G. A. 1994. Size, condition, and territory ownership in male tree swallows (*Tachycineta bicolor*). *Canadian Journal of Zoology* 72:330–333.
- Maan, M. F., T. G. G. Groothuis, and J. Wittenberg. 2001. Escalated fighting despite predictors of conflict outcome: solving the paradox in South American cichlid fish. *Animal Behaviour* 62:623–634.
- Magnhagen, C., and L. Kvarnemo. 1989. Big is better: the importance of size for reproductive success in male *Potamoschisys minutus* (Pallas) (Pisces: Gobiidae). *Journal of Fish Biology* 35:755–763.
- Maynard Smith, J. 1982. *Evolution and the theory of games*. Cambridge University Press, Cambridge.
- Maynard Smith, J., and G. A. Parker. 1976. The logic of asymmetric contests. *Animal Behaviour* 24:159–175.
- Maynard Smith, J., and G. R. Price. 1973. The logic of animal conflict. *Nature* 246:15–18.
- Mayr, M., and A. Berger. 1992. Territoriality and microhabitat selection on two intertidal New Zealand fish. *Journal of Fish Biology* 40:535–544.
- Mesterton-Gibbons, M. 1992. Ecotypic variation in the asymmetric hawk-dove game: when is bourgeois an evolutionarily stable strategy? *Evolutionary Ecology* 6:198–222.
- Mesterton-Gibbons, M., and E. S. Adams. 1998. Animal contests as evolutionary games. *American Scientist* 86:334–341.
- Morrell, L. J., and H. Kokko. 2003. Adaptive strategies of territory formation. *Behavioral Ecology and Sociobiology* 54:385–395.
- . 2005. Bridging the gap between mechanistic and adaptive

- explanations of territory formation. *Behavioral Ecology and Sociobiology* 27:381–390.
- Okada, K., and T. Miyatake. 2004. Sexual dimorphism in mandibles and male aggressive behavior in the presence and absence of females in the beetle *Librodor japonicus* (Coleoptera: Nitidulidae). *Annals of the Entomological Society of America* 97:1342–1346.
- Parker, G. A. 1974. Assessment strategy and the evolution of fighting behaviour. *Journal of Theoretical Biology* 47:223–243.
- Parker, G. A., and D. I. Rubenstein. 1981. Role assessment, reserve strategy, and acquisition of information in asymmetric animal conflicts. *Animal Behaviour* 29:221–240.
- Pereira, H. M., A. Bergman, and J. Roughgarden. 2003. Socially stable territories: the negotiation of space by interacting foragers. *American Naturalist* 161:143–152.
- Petersen, G., and I. C. W. Hardy. 1996. The importance of being larger: parasitoid intruder-owner contests and their implications for clutch size. *Animal Behaviour* 51:1363–1373.
- Pratt, A. E., D. K. McLain, and G. R. Lathrop. 2003. The assessment game in sand fiddler crab contests for breeding burrows. *Animal Behaviour* 65:945–955.
- Pryke, S. R., and S. Andersson. 2003. Carotenoid-based epaulettes reveal male competitive ability: experiments with resident and floater red-shouldered widowbirds. *Animal Behaviour* 66:217–224.
- Ranta, E., and K. Lindström. 1993. Body size and shelter possession in mature signal crayfish, *Pacifasciatus leniusculus*. *Annales Zoologici Fennici* 30:125–132.
- Rosenberg, R. H., and M. Enquist. 1991. Contest behaviour in Weidemeyer's admiral butterfly *Limenitis weidemeyerii* (Nymphalidae): the effect of size and residency. *Animal Behaviour* 42:805–811.
- Stamps, J. A., and V. V. Krishnan. 1999. A learning-based model of territory establishment. *Quarterly Review of Biology* 74:291–318.
- . 2001. How territorial animals compete for divisible space: a learning-based model with unequal competitors. *American Naturalist* 157:154–169.
- Stokkebo, S., and I. C. W. Hardy. 2000. The importance of being gravid: egg load and contest outcome in a parasitoid wasp. *Animal Behaviour* 59:1111–1118.
- Stuart-Smith, A. K., and S. Boutin. 1994. Costs of escalated territorial defence in red squirrels. *Canadian Journal of Zoology* 72:1161–1167.
- Stutt, A. D., and P. Willmer. 1998. Territorial defence in speckled wood butterflies: do the hottest males always win? *Animal Behaviour* 55:1341–1347.
- Switzer, P. V. 2004. Fighting behavior and prior residency advantage in the territorial dragonfly, *Perithemis tenera*. *Ethology Ecology and Evolution* 16:71–89.
- Tobias, J. 1997. Asymmetric territorial contests in the European robin: the role of settlement costs. *Animal Behaviour* 54:9–21.
- Turner, G. F. 1994. The fighting tactics of male mouthbrooding cichlids: the effect of size and residency. *Animal Behaviour* 47:655–662.
- Zack, S., and B. J. Stutchbury. 1992. Delayed breeding in avian social systems: the role of territory quality and floater tactics. *Behaviour* 123:194–219.

Associate Editor: Nicolas Perrin
 Editor: Donald L. DeAngelis